

Anaphylaxis Emergency Action Plan

Patient Name:			Age:	
Allergies:				
Asthma Yes (high risk for sever	re reaction)] No		
Additional health problems besides	s anaphylaxis:			
Concurrent medications:				
Symptoms of Anaphylaxis MOUTH itching, swelling of lips and/or tongue THROAT* itching, tightness/closure, hoarseness SKIN itching, hives, redness, swelling GUT vomiting, diarrhea, cramps LUNG* shortness of breath, cough, wheeze HEART* weak pulse, dizziness, passing out				
		Severity of symptoms life-threatening. ACT		
Emergency Action Steps - D. Inject epinephrine in thigh using (c	OO NOT HESITATE	TO GIVE EPINEPHRI Adrenaclick (0.15 mg)	NE!Adrenaclick (0.3 mg)	
		Auvi-Q (0.15 mg)	☐ Auvi-Q (0.3 mg)	
		EpiPen Jr (0.15 mg)	☐ EpiPen (0.3 mg)	
		nephrine Injection, US (0.15 mg)	P Auto-injector- authorized generic (0.3 mg)	
		Other (0.15 mg)	Other (0.3 mg)	
Specify others:				
IMPORTANT: ASTHMA INHALERS	AND/OR ANTIHIS	TAMINES CAN'T BE D	EPENDED ON IN ANAPHYLAXIS.	
2. Call 911 or rescue squad (before	calling contact)			
3. Emergency contact #1: home		work	cell	
Emergency contact #2: home		work	cell	
Emergency contact #3: home		work	cell	
Comments:				
Doctor's Signature/Date/Phone Numb	per			
Parent's Signature (for individuals ur	nder age 18 yrs)/D	ate		

This information is for general purposes and is not intended to replace the advice of a qualified health professional. For more information, visit www.aaaai.org. © 2017 American Academy of Allergy, Asthma & Immunology 4/2017

